arXiv:0706.3738v2 [math.AG] 9 Jul 2007

EQUIVARIANT LITTLEWOOD-RICHARDSON SKEW TABLEAUX

VICTOR KREIMAN

Abstract. We give a positive equivariant Littlewood-Richardson rule also discovered independently by Molev. Our proof generalizes a proof by Stembridge of the ordinary Littlewood-Richardson rule. We describe a weight-preserving bijection between our indexing tableaux and the Knutson-Tao puzzles.

Contents

1. Introduction 1
2. Statement of Results 2
3. Several Types of Tableaux 8
4. Proofs of Nonnegativity Property and Positivity Criteria 9
5. Generalization of Stembridge's Proof 10
6. Involutions on Reverse Hatted Tableaux 11
7. Proof of Lemma 5.1 14
8. Proof of Lemma 5.2 16
9. Bijection with Knutson-Tao Puzzles 18
References 26

1. Introduction

In [MS], Molev and Sagan introduced a rule in terms of barred tableaux for computing the structure constants $c_{\lambda,\mu}^{\nu}$ for products of two factorial Schur functions. Knutson and Tao [KT] realized that under a suitable specialization these are the structure constants $C_{\lambda,\mu}^{\nu}$ for products of two Schubert classes in the equivariant cohomology ring of the Grassmannian. Knutson and Tao [KT] also gave a new rule for computing $C_{\lambda,\mu}^{\nu}$, i.e., an equivariant Littlewood-Richardson rule, which is manifestly positive in the sense of Graham [Gr]. Their rule was expressed in terms of puzzles, generalizations of combinatorial objects first introduced by Knutson, Tao, and Woodward [KTW].

We describe a new nonnegative equivariant Littlewood-Richardson rule, expressed in terms of skew barred tableaux, which was also discovered independently by Molev [Mo1]. By nonnegative we mean that all of the coefficients are either positive or zero; restricting to the positive coefficients then yields a positive rule. In our proof, we compute the structure constants $c_{\lambda,\mu}^{\nu}$ (as do both [MS] and [Mo1]), and then determine the structure constants $C_{\lambda,\mu}^{\nu}$ by specialization (as does [Mo1]). Our strategy for deriving the structure constants $c_{\lambda,\mu}^{\nu}$ is to generalize a concise proof by

---

Date: February 5, 2008.

Stembridge [St] of the ordinary Littlewood-Richardson rule from Schur functions to factorial Schur functions. This method in fact yields a more general result, namely, a generalization of Zelevinsky's extension of the Littlewood-Richardson rule [Z].

We illustrate a weight-preserving bijction Φ between the skew barred tableaux indexing positive coefficients and the Knutson-Tao puzzles, thus giving a new proof of Knutson and Tao's equivariant Littlewood-Richardson rule, and also demonstrating that our positive rule is really the same rule as Knutson and Tao's, just expressed in terms of different combinatorial indexing sets. We extend Φ to a bijection from all skew barred tableaux indexing nonnegative coefficients to the set of trapezoid puzzles, which are generalizations of puzzles. Our representation of the bijections generalizes Tao's 'proof without words' [V, Figure 11], which gives a bijection between tableaux and puzzles in the nonequivariant setting.

The results of this paper were presented at the AMS Sectional Meeting, Santa Barbara, CA, April 2005, and the University of Georgia Algebra Seminar, August 2006.

2. Statement of Results

Let $\mathbb{N}$ denote the set of nonnegative integers, and let $n \geq d$ be fixed positive integers. For $m \in \mathbb{N}$, define $m' := d + 1 - m$. For $\lambda = (\lambda_1, \ldots, \lambda_d) \in \mathbb{N}^d$, define $|\lambda| = \lambda_1 + \cdots + \lambda_d$. Denote by $\mathcal{P}_d$ the set of all such λ which are **partitions**, i.e., such that $\lambda_1 \geq \cdots \geq \lambda_d$, and by $\mathcal{P}_{d,n}$ the set of all such partitions for which $\lambda_1 \leq n - d$. Let $\lambda = (\lambda_1, \ldots, \lambda_d), \mu = (\mu_1, \ldots, \mu_d), \rho = (d-1, d-2, \ldots, 0)$, and $1 = (1, \ldots, 1)$ be fixed elements of $\mathcal{P}_d$. For any sequence $i = i_1, i_2, \ldots, i_t$, $i_j \in \{1, \ldots, d\}$, define the **content of** i to be $\omega(i) = (\xi_1, \ldots, \xi_d) \in \mathbb{N}^d$, where ξ_k is the number of k's in the sequence.

2.1. Defining the Structure Constants $c^{\nu}_{\lambda,\mu}$ for Products of Factorial Schur Functions.

A **reverse Young diagram** is a right and bottom justified array of boxes. To μ we associate the reverse Young diagram whose bottom row has length μ_1, next to bottom row has length μ_2, etc. We also denote this reverse Young diagram by μ. The columns of a reverse Young diagram are numbered from right to left and the rows from bottom to top.



Figure 1. The reverse Young diagram $(4, 3, 1)$, with rows and columns numbered.

A **reverse tableau of shape** μ is a filling of each box of μ with an integer in $\{1, \ldots, d\}$ in such a way that the entries weakly increase along any row from left to right and strictly increase along any column from top to bottom. Let $\mathcal{R}(\mu)$ denote the set of all reverse tableaux of shape μ. Let $x_1, \ldots, x_d$ be a finite set of variables and $(y_i)_{i\in\mathbb{N}_{>0}}$ an infinite set of variables. For $R \in \mathcal{R}(\mu)$, define

$$(x\,|\,y)^R = \prod_{a\in R} \left(x_a - y_{a'+c(a)-r(a)}\right),$$

where for entry $a \in R$, $c(a)$ and $r(a)$ are the column and row numbers of a respectively. The **factorial Schur function** is defined to be

$$s_\mu(x\,|\,y) = \sum_{R\in\mathcal{R}(\mu)} (x\,|\,y)^R.$$

Factorial Schur functions are special cases of Lascoux and Schützenberger's double Schubert polynomials [LS1, LS2]. Various versions of factorial Schur functions and their properties have been introduced and studied by [BL], [CL], [GG], [La], [Ma1], [Ma2], [Mo1], and [Mo2] (see [Mi], [Mo1], and [MS] for more discussion of these polynomials).

We check that our definition of factorial Schur function agrees with the definition in [MS], which is expressed in terms of Young tableaux. Replacing each entry a in a reverse tableau R by a' and rotating the resulting tableau by 180 degrees, one obtains a Young tableau T. This operation defines a bijection between reverse tableax of shape μ and Young tableaux of shape μ. The polynomials $(x\,|\,y)^T$, as defined in [MS], and $(x\,|\,y)^R$, as defined above, are related by a fixed permutation on the indices of the x_i's, namely the involution $i \mapsto i'$. Thus the equivalence of the two definitions follows from the fact that factorial Schur functions are symmetric in the x_i's. (Corollary 5.4 also establishes the equivalence of the two definitions.)

From the definition of $s_\mu(x\,|\,y)$, one sees that

$$s_\mu(x\,|\,y) = s_\mu(x) + \text{ terms of lower degrees in the } x_i\text{'s},$$

where $s_\mu(x)$ is the Schur function in $x_1, \ldots, x_d$. Since the Schur functions form a $\mathbb{C}$-basis for $\mathbb{C}[x_1, \ldots, x_d]^{S_d}$, the factorial Schur functions must form a $\mathbb{C}[y_i]$-basis for $\mathbb{C}[y_i][x_1, \ldots, x_d]^{S_d}$. Thus

$$s_\lambda(x\,|\,y)s_\mu(x\,|\,y) = \sum c_{\lambda,\mu}^{\nu} s_\nu(x\,|\,y), \tag{1}$$

for some polynomials $c_{\lambda,\mu}^{\nu} \in \mathbb{C}[y_i]$, where the summation is over all $\nu \in \mathcal{P}_d$.

Also from the definition one sees that $s_\mu(x\,|\,y)$ is a homogeneous polynomial of degree $|\mu|$. Therefore $|\lambda| + |\mu| - |\nu| = \deg(c_{\lambda,\mu}^{\nu})$. If $|\lambda| + |\mu| - |\nu| = 0$, then $c_{\lambda,\mu}^{\nu} \in \mathbb{C}$ is the ordinary Littlewood-Richardson coefficient (see [F1], [LR], [Sa]).

2.2. Computing the Structure Constants $c_{\lambda,\mu}^{\nu}$.

The **skew diagram** $\lambda * \mu$ is obtained by placing the Young diagram λ above and to the right of the reverse Young diagram μ (see Figure 2). A **skew barred tableau L of shape $\lambda * \mu$** is a filling of each box of the subdiagram λ of $\lambda * \mu$ with an element of $\{1, \ldots, d\}$ and each box of the subdiagram μ of $\lambda * \mu$ with an element of $\{1, \ldots, d\} \cup \{\overline{1}, \ldots, \overline{d}\}$, in such a way that the values of the entries, without regard to whether or not they are barred, weakly increase along any row from left to right and strictly increase along any column from top to bottom. The **unbarred column word of L**, denoted by L^u, is the sequence of unbarred entries of L beginning at the top of the rightmost column, reading down, then moving to the top of the next to rightmost column and reading down, etc (the barred entries are just skipped over in this process). We say that that the unbarred column word of L is **Yamanouchi** if, when one writes down the word and stops at any point, one will have written at least as many ones as twos, at least as many twos as threes, …, at least as many $(d-1)$'s as d's. The **unbarred content of** L is $\omega(L^u)$, the content of the unbarred column word.

Definition 2.3. *An* ***equivariant Littlewood-Richardson skew tableau*** *is a skew barred tableau whose unbarred column word is Yamanouchi. We denote the set of all equivariant Littlewood-Richardson skew tableaux of shape $\lambda * \mu$ and unbarred content ν by $\mathcal{LR}^{\nu}_{\lambda,\mu}$.*

We remark that this definition forces the i-th row of λ to consist of λ_i unbarred i's.



FIGURE 2. An equivariant Littlewood-Richardson skew tableau of shape $(2,1,1) * (4,3,1)$ and unbarred content $(3,3,2,1)$. The unbarred column word, $1,1,2,3,2,4,3,1,2$, is Yamanouchi, as required.

For L a skew barred tableau and $a \in L$, denote by $L^u_{<a}$ the portion of the unbarred column word of L which comes before reaching a when reading entries from L. Define

$$c_L = \prod_{\substack{a \in L \\ a \text{ barred}}} \left(y_{|a|' + \omega(L^u_{<a})_{|a|}} - y_{|a|' + c(a) - r(a)} \right), \tag{2}$$

where $r(a)$ and $c(a)$ are the row and column numbers of a considered as entries of μ (see Figure 1), and $|a|' = d + 1 - |a|$ (we use the absolute value symbol, $|a|$, to stress that we are interested in the integer *value* of the barred entry a). As usual, the trivial product is defined to be 1. The main result of this paper is the following

Theorem 2.4. $c^{\nu}_{\lambda,\mu} = \sum_{L \in \mathcal{LR}^{\nu}_{\lambda,\mu}} c_L.$

Example 2.5. *Let L be the equivariant Littlewood-Richardson skew tableau of Figure 2. Suppose that $d = 4$. Consider the entry $a = \overline{1}$ in row 2, column 2 of μ. We have $L^u_{<a} = 1,1,2,3,2,4$, so $\omega(L^u_{<a}) = (2,2,1,1)$. Thus $|a|' + \omega(L^u_{<a})_{|a|} = (d+1-(1)) + (2,2,1,1)_1 = 4+2 = 6$. Also, $|a|' + c(a) - r(a) = (d+1-(1)) + 2 - 2 = 4$. Therefore the contribution of this entry to c_L is $y_6 - y_4$.*

Similarly, one computes the contribution of the entry $\overline{2}$ *in row 1, column 3 to be* $y_5 - y_5$ *and the contribution of the entry* $\overline{3}$ *in row 2, column 1 to be* $y_3 - y_1$. *Therefore* $c_L = (y_5 - y_5)(y_6 - y_4)(y_3 - y_1)$, *which equals* 0.

2.6. **Nonnegativity and Positivity.** If $L \in \mathcal{LR}^{\nu}_{\lambda,\mu}$, then we write $c_L > 0$ if each factor in (2) is of the form $y_i - y_j$ with $i > j$. We write $c_L \geq 0$ if either $c_L > 0$ or $c_L = 0$.

Proposition 2.7. *If* $L \in \mathcal{LR}^{\nu}_{\lambda,\mu}$, *then* $c_L \geq 0$.

Let $\mathcal{LR}^{\nu\,+}_{\lambda,\mu}$ be the set of $L \in \mathcal{LR}^{\nu}_{\lambda,\mu}$ for which $c_L > 0$. By Proposition 2.7, we can restrict the summation in Theorem 2.4 to such L:

Corollary 2.8. $c^{\nu}_{\lambda,\mu} = \displaystyle\sum_{L \in \mathcal{LR}^{\nu\,+}_{\lambda,\mu}} c_L.$

One could, of course, use (2), the definition of c_L, to distinguish between $c_L > 0$ and $c_L = 0$: $c_L > 0$ if and only if $\omega(L^{u}_{<a})_{|a|} > c(a) - r(a)$ for all barred $a \in L$. The following Proposition gives a number of other tests for more efficiently making this determination.

Proposition 2.9. *If* $L \in \mathcal{LR}^{\nu}_{\lambda,\mu}$, *then the following are equivalent:*
1. $c_L > 0$
2. $\omega(L^{u}_{<a})_{|a|} > c(a) - r(a)$ *for all barred* $a \in L$.
3. $\omega(L^{u}_{<a})_{|a|} > c(a) - r(a)$ *for all barred* $a \in L$ *with* $r(a) = 1$.
4. $\omega(L^{u}_{<a})_{|a|} \geq c(a)$ *for all barred* $a \in L$
5. $\omega(L^{u}_{<a})_{|a|} \geq c(a)$ *for all barred* $a \in L$ *with* $r(a) = 1$.

If $L \in \mathcal{LR}^{\nu}_{\lambda,\mu}$ satisfies any of these equivalent conditions, then we say that L is **positive**. It is obvious that $4 \implies 2 \implies 3 \iff 5$. Condition 3 states that it suffices to check barred entries on the bottom row of L for positivity. Condition 4 has the following interpretation: for any barred entry $a \in L$, the corresponding factor $y_i - y_j$ in c_L satisfies $i - j \geq r(a)$ (which of course implies $i - j > 0$, the condition required for positivity).

Example 2.10. *Let* $d = 3$, $\lambda = (1,1)$, $\mu = (3,2)$, *and* $\nu = (3,2,1)$. *We list all* $L \in \mathcal{LR}^{\nu\,+}_{\lambda,\mu}$, *and for each* L *we give* c_L:



$c_L = y_6 - y_5$



$c_L = y_5 - y_3$



$c_L = y_4 - y_2$



$c_L = y_3 - y_1$

Note that if L *has an unbarred* 2 *in the upper right box of* μ, *then the unbarred column word of* L *is not Yamanouchi, and if* L *has two unbarred* 1*'s on the top row of* μ *and is not the leftmost diagram, then* $c_L = 0$; *thus we do not include such* L *among* $\mathcal{LR}^{\nu\,+}_{\lambda,\mu}$. *By Corollary 2.8,* $c^{\nu}_{\lambda,\mu} = (y_6 - y_5) + (y_5 - y_3) + (y_4 - y_2) + (y_3 - y_1) = (y_6 - y_1) + (y_4 - y_2)$.

We list all $L \in \mathcal{LR}^{\nu\,+}_{\mu,\lambda}$, *and for each* L *we give* c_L:



$c_L = y_6 - y_2$



$c_L = y_4 - y_1$

By Corollary 2.8, $c^{\nu}_{\mu,\lambda} = (y_6 - y_2) + (y_4 - y_1)$*. We see that* $c^{\nu}_{\mu,\lambda} = c^{\nu}_{\lambda,\mu}$*. This is a general fact ensured by (1); however, it is not apparent from the statement of Corollary 2.8.*

Example 2.11. *For cases where* $\mu = \nu$*, a formula for* $c^{\nu}_{\lambda,\mu}$ *which produces a different positive expression than Corollary 2.8 appears in* [Bi], [IN]*, and* [Kr]*. For example, using this formula, for* $d = 3$*,* $\lambda = (2,1)$*, and* $\mu = \nu = (3,3,1)$*, one computes:*

$$c^{\nu}_{\lambda,\mu} = (y_6 - y_1)(y_6 - y_3)(y_5 - y_1) + (y_6 - y_1)(y_5 - y_4)(y_5 - y_1).$$

Using Corollary 2.8:

$$\begin{aligned} c^{\nu}_{\lambda,\mu} =& (y_5 - y_3)(y_5 - y_1)(y_3 - y_1) + (y_6 - y_4)(y_5 - y_1)(y_3 - y_1) \\ &+ (y_6 - y_4)(y_6 - y_3)(y_3 - y_1) + (y_5 - y_3)(y_4 - y_3)(y_5 - y_1) \\ &+ (y_6 - y_4)(y_4 - y_3)(y_5 - y_1) + (y_6 - y_4)(y_6 - y_3)(y_4 - y_3) \\ &+ (y_6 - y_4)(y_5 - y_4)(y_5 - y_1) + (y_6 - y_4)(y_5 - y_4)(y_6 - y_3) \end{aligned}$$

$$\begin{aligned} c^{\nu}_{\mu,\lambda} =& (y_6 - y_4)(y_6 - y_2)(y_5 - y_2) + (y_5 - y_3)(y_6 - y_2)(y_5 - y_2) \\ &+ (y_6 - y_4)(y_6 - y_2)(y_2 - y_1) + (y_5 - y_3)(y_6 - y_2)(y_2 - y_1) \\ &+ (y_6 - y_4)(y_5 - y_1)(y_2 - y_1) + (y_5 - y_3)(y_5 - y_1)(y_2 - y_1). \end{aligned}$$

These three polynomials are, of course, equal.

For $L \in \mathcal{LR}^{\nu}_{\lambda,\mu}$, $|\lambda| + |\mu| - |\nu| = \#(\text{entries of } L) - \#(\text{unbarred entries of } L) = \#(\text{barred entries of } L) = \deg(c^{\nu}_{\lambda,\mu})$. In particular, if $|\lambda| + |\mu| - |\nu| = 0$, then L has no barred entries. When $|\lambda| + |\mu| - |\nu| = 0$, Theorem 2.4 is the ordinary Littlewood-Richardson rule (see [F1], [LR], [Sa]).

2.12. **Defining the Structure Constants $C^{\nu}_{\lambda,\mu}$ for products of two Schubert Classes in $H^*_T(Gr_{d,n})$.** The **Grassmannian** $Gr_{d,n}$ is the set of d-dimensional complex subspaces of $\mathbb{C}^n$. Let $\{e_1, \dots, e_n\}$ be the standard basis for $\mathbb{C}^n$. Consider the opposite standard flag, whose i-th space is $\mathrm{Span}(e_n, \dots, e_{n-i+1})$. For $\lambda \in \mathcal{P}_{d,n}$, the (opposite) Schubert variety X_λ of $Gr_{d,n}$ is defined by incident relations:

$$X_\lambda = \{V \in Gr_{d,n} \mid \dim(V \cap F_i) \geq \dim(\mathbb{C}^\lambda \cap F_i)\},\ i = 1, \dots, n,$$

where $C^\lambda = \mathrm{Span}(e_{\lambda_d+d}, \dots, e_{\lambda_1+1})$. The Schubert variety X_λ is invariant under the action of the group $T = (\mathbb{C}^*)^n$ on $Gr_{d,n}$. Thus it determines a class S_λ in the equivariant cohomology ring $H^*_T(Gr_{d,n})$.

Let $V = Gr_{d,n} \times \mathbb{C}^n$ be the trivial vector bundle on $Gr_{d,n}$, with diagonal T-action, where T acts naturally on $Gr_{d,n}$ and on $\mathbb{C}^n$ (thus V is not equivariantly trivial). Let $S = \{(w,v) \in V \mid v \in w\}$ be the tautological vector bundle on $Gr_{d,n}$. Then S is a T-invariant sub-bundle of V. Let $Y_1, \dots, Y_n$ be the equivariant Chern roots of V^* and $X_1, \ldots, X_d$ the equivariant Chern roots of S. Then $H_T^*(Gr_{d,n})$ is a free $\mathbb{C}[Y_1, \ldots, Y_n]$-module, with the Schubert classes forming a $\mathbb{C}[Y_1, \ldots, Y_n]$-basis. Thus for $\lambda, \mu \in \mathcal{P}_{d,n}$,

$$S_\lambda S_\mu = \sum C_{\lambda,\mu}^{\nu} S_\nu,$$

for some $C_{\lambda,\mu}^{\nu} \in \mathbb{C}[Y_1, \ldots, Y_n]$, where the summation is over all $\nu \in \mathcal{P}_{d,n}$. We have (see [F2], [KT], [Mi])

Proposition 2.13. *For* $\lambda \in \mathcal{P}_d$, $S_\lambda = s_\lambda(X_1, \ldots, X_d, -Y_n, \ldots, -Y_1, 0, 0, \ldots)$.

Thus by specializing (1), we can determine the structure constants $C_{\lambda,\mu}^{\nu}$.

Corollary 2.14. *For* $\lambda, \mu, \nu \in \mathcal{P}_{d,n}$, $C_{\lambda,\mu}^{\nu} = c_{\lambda,\mu}^{\nu}(-Y_n, \ldots, -Y_1, 0, 0, \ldots)$.

2.15. **Computing the Structure Constants $C_{\lambda,\mu}^{\nu}$.** Let $\lambda, \mu, \nu \in \mathcal{P}_{d,n}$. By Corollary 2.14, the structure constant $C_{\lambda,\mu}^{\nu}$ can be computed using the formula for $c_{\lambda,\mu}^{\nu}$. Let $L \in \mathcal{LR}_{\lambda,\mu}^{\nu}$. Define

$$\begin{aligned} C_L &= c_L(-Y_n, \ldots, -Y_1) \\ &= \prod_{\substack{a \in L \\ a \text{ barred}}} \left(Y_{(n-d)+|a|-(c(a)-r(a))} - Y_{(n-d)+|a|-\omega(L^u_{<a})_{|a|}}\right). \end{aligned} \tag{3}$$

We write $C_L > 0$ if each factor in (3) is of the form $Y_i - Y_j$ with $i > j$, and we write $c_L \geq 0$ if either $C_L > 0$ or $C_L = 0$. By (3), $c_L \geq 0 \iff C_L \geq 0$, and $c_L = 0 \iff C_L = 0$. Thus Propositions 2.7 and 2.9 imply

Corollary 2.16. $C_L \geq 0$*, and* $C_L > 0 \iff L$ *satisfies any of the equivalent conditions of Proposition 2.9.*

By Theorem 2.4, Corollary 2.14, and Corollary 2.16, we have

Corollary 2.17. $C_{\lambda,\mu}^{\nu} = \displaystyle\sum_{L \in \mathcal{LR}_{\lambda,\mu}^{\nu}} C_L = \sum_{L \in \mathcal{LR}_{\lambda,\mu}^{\nu\,+}} C_L.$

Example 2.18. *We continue Example 2.10. For* $n \geq 6$, $\lambda, \mu \in \mathcal{P}_{d,n}$. *Thus for* $\nu \in \mathcal{P}_{d,n}$, $C_{\lambda,\mu}^{\nu} = (Y_{n+1-2} - Y_{n+1-6}) + (Y_{n+1-1} - Y_{n+1-4}) = (Y_{n-1} - Y_{n-5}) + (Y_n - Y_{n-3})$.

2.19. **Equivalence of Molev's Results.** Our equivariant Littlewood-Richardson skew tableaux are essentially the same as the Molev's indexing tableaux [Mo1]. To determine the tableau in [Mo1] which corresponds to our $L \in \mathcal{LR}_{\lambda,\mu}^{\nu}$, replace all barred entries of $L|_\mu$ by unbarred entries and visa-versa, and then rotate the resulting object by 180 degrees. If one makes this modification, then Corollary 2.8 is equivalent to [Mo1, Theorem 2.1] after accounting for the relationship between double Schur polynomials and factorial Schur functions (see [Mo1, (1.9)]), and Corollary 2.17 is identical to [Mo1, Corollary 3.1].

In our notation, Molev's positivity criterion states that for $L \in \mathcal{LR}_{\lambda,\mu}^{\nu}$, $c_L > 0$ if and only if

$$\omega(L^u)'_{c(a)} \geq |a| \text{ for all } a \in L \text{ with } r(a) = 1, \tag{4}$$

where $\omega(L^u)'$ is the *conjugate partition* to $\omega(L^u)$ (in this case Molev calls $L|_\mu$ ν-*bounded*). One can re-express (4) as follows:

$$\omega(L^u)_{|a|} \geq c(a) \text{ for all } a \in L \text{ with } r(a) = 1.$$

It is not difficult to see that this condition is equivalent to Proposition 2.9.5.

Related and more general results have been achieved in several directions. Robinson [R] has given a Pieri rule in the equivariant cohomology of the flag variety. McNamara [Mc] introduced factorial Grothendieck polynomials, generalizations of factorial Schur functions, and has given a rule for computing the structure constants for various of their products.

This paper is organized as follows. In Section 3, we introduce various types of tableaux which will appear throughout the paper. In Section 4, we prove Propositions 2.7 and 2.9, the nonnegativity property and positivity criteria of c_L. In Section 5, we outline the main steps in our proof of Theorem 2.4, whose two difficult technical lemmas are proved in Sections 7 and 8. In Section 6, we define a set of involutions required for the proofs of these two lemmas. In Section 9, we describe a bijection between positive equivariant Littlewood-Richardson skew tableaux and Knutson-Tao puzzles.

3. Several Types of Tableaux

In this section we collect the definitions of the several types of tableaux which we will encounter in the remainder of the paper: reverse barred tableaux, reverse barred subtableaux, and reverse hatted tableaux. The latter two are refinements of the first.

A **reverse barred tableau of shape** μ is a skew barred tableau of shape $\emptyset * \mu$; alternatively, it can be defined as a reverse Young diagram of shape μ, each of whose boxes is filled with either an integer k or a barred integer $\overline{k}$, $k \in \{1, \ldots, d\}$, in such a way that the values of the entries, without regard to whether or not they are barred, weakly increase along any row from left to right and strictly increase along any column from top to bottom. We denote the set of all reverse barred tableaux of shape μ by $\mathcal{B}(\mu)$. If $B \in \mathcal{B}(\mu)$, then define $\lambda * B$ to be the skew barred tableau obtained by placing the Young tableau whose i-th row consists of λ_i i's above and to the right of B. Then $B \mapsto \lambda * B$ defines a bijection from $\{B \in \mathcal{B}(\mu) \mid (\lambda * B)^u$ is Yamanouchi$\}$ to the equivariant Littlewood-Richardson skew tableaux of shape $\lambda * \mu$, whose inverse map is $L \mapsto L|_\mu$. Any $a \in B$ also corresponds to an entry $a \in \lambda * B$. Define B^u and $B^u_{<a}$ to be $(\emptyset * B)^u$ and $(\emptyset * B)^u_{<a}$ respectively.

A **reverse barred subtableaux of shape** μ is a reverse Young diagram μ each of whose boxes contains either an integer k, a barred integer $\overline{k}$, or is empty, where $k \in \{1, \ldots, d\}$. A **reverse subtableau of shape** μ is a reverse barred tableau of shape μ which has no barred entries. We do not define any notion of row semistrictness or column strictness for such objects, as no such conditions will be required for our purposes. Denote the set of all reverse subtableaux and reverse barred subtableaux of shape μ by $\mathcal{R}_{sub}(\mu)$ and $\mathcal{B}_{sub}(\mu)$ respectively. We have the following containments:

$$\begin{array}{ccc}
\mathcal{R}_{sub}(\mu) & \subset & \mathcal{B}_{sub}(\mu) \\
\cup & & \cup \\
\mathcal{R}(\mu) & \subset & \mathcal{B}(\mu)
\end{array}$$

For $B \in \mathcal{B}_{sub}(\mu)$ and $a \in B$, define B^u and $B^u_{<a}$ just as for elements of $\mathcal{B}(\mu)$, assuming that when reading the unbarred column word of B, both barred entries and empty boxes are skipped over. If $B \in \mathcal{B}_{sub}(\mu)$, then define $\widetilde{B} \in \mathcal{R}_{sub}(\mu)$ to be the reverse subtableau obtained by removing all bars from entries of B, i.e., replacing each barred entry of B by an unbarred entry of the same value.

A **reverse hatted tableau of shape** μ is a reverse Young diagram μ each of whose boxes is filled with either a(n) (un-hatted) integer k, a *left* hatted integer $\check{k}$, or a *right* hatted integer $\hat{k}$, $k \in \{1, \ldots, d\}$, such that the values of the entries, without regard to whether or not they are hatted, weakly increase along any row from left to right and strictly increase along any column from top to bottom. Denote the set of all reverse hatted tableaux of shape μ by $\mathcal{H}(\mu)$. If H is a reverse hatted tableau, then define $\overline{H}$ to be the reverse barred tableau produced by replacing all hats (right and left) by bars. Hence for a reverse barred tableau B with m barred entries, there are 2^m reverse hatted tableaux H such that $\overline{H} = B$ (since each $\overline{k}$ of B can be replaced by either $\check{k}$ or $\hat{k}$). For $a \in H$, define H^u and $H^u_{<a}$ to be $\overline{H}^u$ and $\overline{H}^u_{<a}$ respectively. Define H^l (resp. H^r) to be the set of left-hatted (resp. right-hatted) entries of H.

We next give two different ways to generalize the polynomial c_L defined in Section 2. Let $\xi \in \mathbb{N}^d$. For $B \in \mathcal{B}_{sub}(\mu)$, define

$$c_{\xi,B} = \prod_{\substack{a \in B \\ a \text{ barred}}} (y_{e_{\xi,B}(a)} - y_{f_B(a)}), \tag{5}$$

where $e_{\xi,B}(a) := (\xi + \omega(B^u_{<a}))_{|a|}$ and $f_B(a) := |a|' + c(a) - r(a)$, $a \in B$. For $H \in \mathcal{H}(\mu)$, define

$$d_{\xi,H} = \prod_{a \in H^l} y_{e_{\xi,H}(a)} \prod_{a \in H^r} (-y_{f_H(a)}), \tag{6}$$

where $e_{\xi,H}(a) := (\xi + \omega(H^u_{<a}))_{|a|}$ and $f_H(a) := |a|' + c(a) - r(a)$, $a \in H$. In both (5) and (6), the empty product is defined to equal 1.

Let $B \in \mathcal{B}(\mu)$. By definition,

$$c_{\lambda * B} = c_{\lambda+\rho+1,B}. \tag{7}$$

In addition, the equation

$$c_{\xi,B} = \sum_{\substack{H \in \mathcal{H}(\mu) \\ \overline{H} = B}} d_{\xi,H} \tag{8}$$

expresses $c_{\xi,B}$ by expanding (5) in terms of monomials in the y_i's. Combining (7) and (8), we have

$$c_{\lambda * B} = \sum_{\substack{H \in \mathcal{H}(\mu) \\ \overline{H} = B}} d_{\lambda+\rho+1,H}. \tag{9}$$

If $R \in \mathcal{R}_{sub}(\mu)$, then define $(x \,|\, y)^R = \prod_{a \in R} \big(x_a - y_{f_R(a)}\big)$. This definition is consistent with the definition of $(x \,|\, y)^R$, $R \in \mathcal{R}(\mu)$, given in Section 2.

4. PROOFS OF NONNEGATIVITY PROPERTY AND POSITIVITY CRITERIA

Let $L \in \mathcal{LR}^{\nu}_{\lambda,\mu}$, and let $B = L|_{\mu}$. For $a \in B$, which we also view as an entry of L, define $L^u_{\leq a}$ to be $L^u_{<a}$ if a is barred, or $L^u_{<a}$ appended with a if a is not barred. Define

$$\Delta(a) = \omega(L^u_{\leq a})_{|a|} - c(a) + r(a).$$

If a is barred, then $\omega(L^u_{\leq a}) = \omega(L^u_{<a})$; hence $\Delta(a)$ gives the difference between the two indices $i - j$ of the factor $y_i - y_j$ corresponding to a in (2). Therefore Propositions 2.7 and 2.9 are equivalent to the following two lemmas respectively.

Lemma 4.1. *If* $\Delta(a) < 0$ *for some barred* $a \in B$*, then* $\Delta(b) = 0$ *for some barred* $b \in B$*.*

Lemma 4.2. *The following are equivalent:*
(i) $\Delta(a) > 0$ *for all barred* $a \in B$*.*
(ii) $\Delta(a) > 0$ *for all barred* $a \in B$ *with* $r(a) = 1$*.*
(iii) $\Delta(a) \geq r(a)$ *for all barred* $a \in B$*.*

Before proving these two lemmas, we first establish some properties of Δ.

Lemma 4.3. *The function* $\Delta : B \to \mathbb{Z}$ *satisfies the following properties:*
(i) If $a \in B$ *and* $c(a) = 1$*, then* $\Delta(a) \geq 0$*, with equality implying that* a *is barred.*
(ii) If one moves left by one box, then Δ *can decrease by at most one. If it does decrease by one, then the left box must be barred.*
(iii) If $\Delta(a) \leq 0$ *for some* $a \in B$*, then* $\Delta(b) = 0$ *for some barred* $b \in B$ *on the same row as* a*.*
(iv) The function $a \mapsto \Delta(a) - r(a)$ *is weakly decreasing as one moves down along any column.*

Proof. (i) Since $r(a) \geq 1$, $\Delta(a) \geq 0$. If $\Delta(a) = 0$, then $r(a) = 1$ and $\omega(L^u_{\leq a})_{|a|} = 0$. The latter requirement implies that a is barred.
(ii) If entry m lies one box left of a, then $-c(m) = -c(a) - 1$, $r(m) = r(a)$, and $\omega(L^u_{\leq m})_{|m|} \geq \omega(L^u_{\leq a})_{|m|} \geq \omega(L^u_{\leq a})_{|a|}$, where the first inequality is an equality if and only if m is barred. The second inequality is a consequence of the fact that the unbarred column word of L is Yamanouchi.
(iii) Let m be rightmost entry in the same row as a. If $\Delta(m) = 0$, then by (i), m is barred, so letting $b = m$ we are done. Otherwise $\Delta(m) > 0$. By (ii), as one moves left from m to a along the row the two entries lie on, one must encounter some barred b for which $\Delta(b) = 0$.
(iv) If entry m lies one box below a, then $\omega(L^u_{\leq a})_{|a|} = \omega(L^u_{\leq m})_{|a|} \geq \omega(L^u_{\leq m})_{|m|}$, since the unbarred column word of L is Yamanouchi. ☐

Proof of Lemmas 4.1 and 4.2. Lemma 4.1 is a special case of Lemma 4.3(iii). In Lemma 4.2, implications (iii) $\Longrightarrow$ (i) $\Longrightarrow$ (ii) are clear. We prove (ii) $\Longrightarrow$ (iii). Suppose that $a \in B$ is a barred entry such that $\Delta(a) < r(a)$. Let m be the bottom entry in column $c(a)$. By Lemma 4.3(iv), $\Delta(m) < r(m)$. Since $r(m) = 1$, $\Delta(m) \leq 0$. By Lemma 4.3(iii), $\Delta(b) = 0$ for some barred b on the bottom row of B. ☐

5. Generalization of Stembridge's Proof

In this section we list the main steps in the proof of Theorem 2.4. The bulk of the technical work, however, namely the proofs of Lemmas 5.1 and 5.2, is taken up in the three subsequent sections. The underlying logic and structure of our arguments in this and the following three sections follows Stembridge [St], who works out similar results for ordinary Schur functions.

For $k \in \mathbb{N}$, define the polynomial $(x_j \,|\, y)^k = (x_j - y_1) \cdots (x_j - y_k)$. For $\xi = (\xi_1, \ldots, \xi_d) \in \mathbb{N}^d$, define $a_\xi(x \,|\, y) = \det[(x_j \,|\, y)^{\xi_i}]_{1 \leq i,j \leq d}$.

Lemma 5.1. $a_{\lambda+\rho}(x \,|\, y) s_\mu(x \,|\, y) = \displaystyle\sum_{B \in \mathcal{B}(\mu)} c_{\lambda * B}\, a_{\lambda+\rho+\omega(B^u)}(x \,|\, y).$

Lemma 5.2. *$\sum c_{\lambda * B} a_{\lambda+\rho+\omega(B^u)}(x \,|\, y) = 0$, where the sum is over all $B \in \mathcal{B}(\mu)$ such that the unbarred column word of $\lambda * B$ is not Yamanouchi.*

The following four corollaries follow easily from these two lemmas.

Corollary 5.3. *$a_{\lambda+\rho}(x \,|\, y) s_\mu(x \,|\, y) = \sum c_{\lambda * B} a_{\lambda+\rho+\omega(B^u)}(x \,|\, y)$, where the sum is over all $B \in \mathcal{B}(\mu)$ such that the unbarred column word of $\lambda * B$ is Yamanouchi.*

Suppose that $B \in \mathcal{B}(\mu)$ is such that the unbarred column word of $\emptyset * B$ is Yamanouchi. If B has barred entries, then by Propositions 2.7 and 2.9.5, $c_{\emptyset * B} = 0$. If B has no barred entries, then B must be the unique reverse tableau of shape μ and content μ: B contains a 1 at the top of each column, and its entries increase by 1 per box as one moves down any column. Thus, by setting $\lambda = \emptyset$ in Corollary 5.3, we arrive at a new proof of the bialternant formula for the factorial Schur function ([GG], [Ma1]):

Corollary 5.4. *$s_\mu(x \,|\, y) = a_{\mu+\rho}(x \,|\, y)/a_\rho(x \,|\, y)$.*

Dividing both sides of the equation in Corollary 5.3 by $a_\rho(x \,|\, y)$ and applying Corollary 5.4 yields

Corollary 5.5. *$s_\lambda(x \,|\, y) s_\mu(x \,|\, y) = \sum c_{\lambda * B} s_{\lambda+\omega(B^u)}(x \,|\, y)$, where the sum is over all $B \in \mathcal{B}(\mu)$ such that the unbarred column word of $\lambda * B$ is Yamanouchi.*

Regrouping the terms in this summation:

$$s_\lambda(x \,|\, y) s_\mu(x \,|\, y) = \sum_\nu \left(\sum_{\substack{B \in \mathcal{B}(\mu) \\ (\lambda * B)^u \text{ Yamanouchi} \\ \lambda+\omega(B^u)=\nu}} c_{\lambda * B} \right) s_\nu(x \,|\, y) = \sum_\nu \left(\sum_{L \in \mathcal{LR}^\nu_{\lambda,\mu}} c_L \right) s_\nu(x \,|\, y).$$

This proves Theorem 2.4.

Remark 5.6. *Let $\kappa \in \mathcal{P}_d$, $\kappa \le \mu$, i.e., $\kappa_i \le \mu_i$, $i = 1, \dots, d$. One can extend our analysis to factorial skew Schur functions of the form $s_{\mu/\kappa}(x \,|\, y)$ (see* [Ma1]*). One replaces $\mathcal{B}(\mu)$ with $\mathcal{B}(\mu/\kappa)$, the set of all reverse barred tableaux of shape μ/κ. All above definitions extend naturally. For example, for $B \in \mathcal{B}(\mu/\kappa)$, $c_{\lambda * B}$ is computed just as for $B \in \mathcal{B}(\mu)$, but with all boxes of $\kappa \subset \mu$ assumed to be empty. All proofs are virtually unchanged, modified only by formally replacing μ by μ/κ. As a generalization of Corollary 5.5, we obtain*

$$s_\lambda(x \,|\, y) s_{\mu/\kappa}(x \,|\, y) = \sum c_{\lambda * B} s_{\lambda+\omega(B^u)}(x \,|\, y),$$

where the sum is over all $B \in \mathcal{B}(\mu/\kappa)$ such that $(\lambda * B)^u$ is Yamnaouchi. This generalizes Zelevinsky's extension of the Littlewood-Richardson rule ([St], [Z]*).*

6. Involutions on Reverse Hatted Tableaux

In his proof, Stembridge [St] utilizes involutions on Young tableaux introduced by Bender and Knuth [BK]. There is an analogous set of involutions on $\mathcal{H}(\mu)$ which satisfy properties required for the proofs of Lemmas 5.1 and 5.2 (see Lemma 6.4). We remark that we were unable to find a suitable set of involutions on $\mathcal{B}(\mu)$, and this is what initially led us to examine $\mathcal{H}(\mu)$. If the involutions on $\mathcal{H}(\mu)$ are restricted to $\mathcal{R}(\mu)$, then the Bender-Knuth involutions are recovered.

6.1. **The Involutions $\mathbf{s_1}, \ldots, \mathbf{s_{d-1}}$ of $\mathcal{H}(\mu)$.** Let $H \in \mathcal{H}(\mu)$, and let $i \in \{1, \ldots, d-1\}$ be fixed. Then an entry a of H with value i or $i+1$ is

- **free** if there is no entry of value $i+1$ or i respectively in the same column;
- **semi-free** if there is an entry of value $i+1$ or i respectively in the same column, and at least one of the two is hatted; or
- **locked** if there is an entry of value $i+1$ or i respectively in the same column, and both entries are unhatted.

Note that any entry of value i or $i+1$ must be exactly one of these three types, and each hatted entry of value i or $i+1$ must be either free or semi-free. In any row, the free entries are consecutive. Semi-free entries come in pairs, one below the other, as do locked entries.

To define the action of s_i on $H \in \mathcal{H}(\mu)$, we first consider how it modifies the free entries of H (see Example 6.2):

1. Let S be a maximal string of free entries with values i and $i+1$ on some row of H. Let S^0, S^l, and S^r denote the unhatted, left-hatted, and right-hatted entries of S respectively. Modify $S^\circ \cup S^l$, as follows:
   A Change the value of each entry of value i to $i+1$ and each entry of value $i+1$ to i, without changing whether or not it has a left hat.
   B Swap the entries of value i with those of value $i+1$: remove all entries of value i; then move each entry of value $i+1$, beginning with the rightmost one, into the rightmost available empty box; then put the removed entries of value i back into the empty boxes of B, preserving the relative order of barred and unbarred entries.

   In this step, $S^\circ \cup S^l$ has been modified. No other entries of H, in particular no entries of S^r, have been modified, changed, or moved. Denote the modified string S by S_1. A potential problem has been introduced: the values of the entries of S_1 may not be weakly increasing as one moves from left to right. In step 2 we correct for this.
2. Let $(S_1^r)_i$ and $(S_1^r)_{i+1}$ denote the entries of S_1^r of value i and $i+1$ respectively. Beginning with the leftmost entry $a \in (S_1^r)_i$, let b be the entry of S_1 to the left of a. If b has value $i+1$, then switch the entries b and a, and then change the left entry from $\widehat{i}$ to $\widehat{i+1}$. Now move right to the next entry of $(S_1^r)_i$, and repeat this procedure until it has been performed on all entries of $(S_1^r)_i$. Next, beginning with the rightmost entry $a \in (S_1^r)_{i+1}$, let b be the entry of S_1 to the right of a. If b has value i, then switch the entries b and a, and then change the right entry from $\widehat{i+1}$ to $\widehat{i}$. Now move left to the next entry of $(S_1^r)_{i+1}$, and repeat this procedure until it has been performed on all entries of $(S_1^r)_{i+1}$.

   Upon completion, we denote by S_2 the resulting string obtained by modifying S_1. It is weakly increasing.

We next consider how s_i modifies the semi-free entries of H:

3. For a semi-free pair consisting of two entries lying in the same column of H, each entry removes its hat (if it has one) and places it on top of the other entry.

The reverse tableau s_iH is obtained by applying steps 1 and 2 to each maximal string S of free entries of H (replacing S by S_2) and then applying step 3 to each semi-free pair.

Example 6.2. *We illustrate steps 1 and 2. Suppose that $i = 2$, and S consists of the following maximal string of consecutive free entries lying along some row of H:*

$$\begin{array}{rcccccccccccc}
S = & 2 & \hat{2} & \check{2} & 2 & 2 & \check{2} & \hat{2} & 2 & 3 & \check{3} & \hat{3} & \check{3} \\
S^0 \cup S^l = & 2 & & \check{2} & 2 & 2 & \check{2} & & 2 & 3 & \check{3} & & \check{3} \\
 & 3 & & \check{3} & 3 & 3 & \check{3} & & 3 & 2 & \check{2} & & \check{2} \\
 & 2 & & \check{2} & \check{2} & 3 & \check{3} & & 3 & 3 & \check{3} & & 3 \\
S_1 = & 2 & \hat{2} & \check{2} & \check{2} & 3 & \check{3} & \hat{2} & 3 & 3 & \check{3} & \hat{3} & 3 \\
S_2 = & 2 & \hat{2} & \check{2} & \check{2} & 3 & \hat{3} & \check{3} & 3 & 3 & \check{3} & \hat{3} & 3
\end{array}$$

In line 2 we remove the entries of S^r from the picture for convenience, in order to focus attention on the operations performed in step 1, which only affect $S^0 \cup S^l$. In lines 3 and 4 the results of applying steps 1A and 1B successively to $S^0 \cup S^l$ are shown. In line 5, the removed entries from S^r are replaced. In line 6, the result of applying step 2 to S_1 is shown. Only two entries are changed in this step.

This algorithm defines maps $b_l : H^l \to (s_iH)^l$ and $b_r : H^r \to (s_iH)^r$, as follows. If $a \in H^l$ is free, then in step 1A, the value of a is either increased or decreased by 1; in step 1B, it is then moved to a different box; in step 2, this new entry in this new box is moved at most one box and changed by at most one in value, resulting in the entry we denote by $b_l(a)$. If $a \in H^r$ is free, then a is unchanged in step 1 and moved at most one box and changed by at most one in value in step 2. Denote the resulting entry by $b_r(a)$. If $a \in H^l$ or $a \in H^r$ is semi-free, then $b_l(a)$ or $b_r(a)$ is the entry in s_iH which it gives its hat to.

In Example 6.2, if a is the rightmost entry of S, which is a $\check{3}$, then $b_l(a)$ is the $\check{2}$ which is the fourth entry of S_2 from left. These two entries are, of course, entries of H and s_iH respectively.

Lemma 6.3. *s_i is an involution on $\mathcal{H}(\mu)$, $i \in \{1, \dots, d-1\}$.*

Proof. We begin by showing that $s_iH \in \mathcal{H}(\mu)$, i.e., s_iH is row semistrict and column strict. The only nonobvious condition is that if S is any maximal string of free entries of H lying along some row, and S_2 the string that replaces it in s_iH, then s_iH weakly increases along the left and right boundaries of S_2. To see this, note that if any entry of H of value $i+1$ is free, then so are all entries of value $i+1$ to the right of it in the same row; and if any entry of H of value i is free, then so are all entries of value i to the left of it in the same row. Thus by the maximality of S, there are no entries of H of value i in the same row and to the right of S, and there are no entries of H of value $i+1$ in the same row and to the left of S. Hence changing values of S from i to $i+1$ and visa-versa to form S_2 does not affect the row semistrictness of H along its boundaries.

We next show that $s_i^2 = \mathrm{id}$. Since the free entries of H lie in the same boxes as the free entries of s_iH, it suffices to show that $s_i^2(S) = S$ for any maximal string S of free entries of H (where s_iS is defined to be s_iH restricted to S). If step 1 is applied to $(s_iS)^\circ \cup (s_iS)^l$, then one sees that the same entries of $S^\circ \cup S^l$ are retrieved, although possibly not in their same boxes. However the relative order of the entries is the same. Now one checks that for $a \in H^r$, $b_r^2(a) = a$. □

Let σ_i be the simple transposition of the permutation group S_d which exchanges i and $i+1$. The involution s_i satisfies the following properties:

Lemma 6.4. *Let $H \in \mathcal{H}(\mu)$, $a \in H^l$, and $b \in H^r$. Then*
(i) $|b_l(a)| = \sigma_i |a|$
(ii) $\omega((s_i H)^u) = \sigma_i \omega(H^u)$.
(iii) $\omega((s_i H)^u_{<b_l(a)}) = \sigma_i \omega(H^u_{<a})$
(iv) $e_{\sigma_i \xi, s_i H}(b_l(a)) = e_{\xi,H}(a)$
(v) $f_{s_i H}(b_r(b)) = f_H(b)$
(vi) $d_{\sigma_i \xi, s_i H} = d_{\xi,H}$

Proof. (i), (ii), and (iii) follow from the construction of s_i.
(iv) By parts (iii) and (i),

$$\begin{aligned} e_{\sigma_i\xi,s_iH}(b_l(a)) &= (\sigma_i\xi + \omega((s_iH)^u_{<b_l(a)}))_{|b_l(a)|} \\ &= (\sigma_i\xi + \sigma_i\omega(H^u_{<a}))_{|b_l(a)|} \\ &= (\sigma_i\xi + \sigma_i\omega(H^u_{<a}))_{\sigma_i|a|} \\ &= (\sigma_i(\xi + \omega(H^u_{<a})))_{\sigma_i|a|} = (\xi + \omega(H^u_{<a}))_{|a|} = e_{\xi,H}(a) \end{aligned}$$

(v) Under b_r, the entry b is either kept in place, moved up, down, left, or right by one box. In these cases, its value is either left unchanged, decreased, increased, increased, or decreased by one respectively. The result now follows from the definition of f_H.
(vi) This is a consequence of (i), (ii), and (6). □

Let $H \in \mathcal{H}(\mu)$ and let $\sigma \in S_d$. Choose some decomposition of σ into simple transpositions: $\sigma = \sigma_{i_1} \cdots \sigma_{i_t}$. Define $\sigma H := s_{i_1} \cdots s_{i_t} H$. Although σH depends on the decomposition chosen for σ, by Lemma 6.4(ii) and (vi),

$$\omega((\sigma H)^u) = \sigma\omega(H^u) \qquad \text{and} \qquad d_{\sigma\xi,\sigma H} = d_{\xi,H}. \tag{10}$$

In particular, both $\omega((\sigma H)^u)$ and $d_{\sigma\xi,\sigma H}$ are independent of the decomposition of σ.

7. PROOF OF LEMMA 5.1

Lemma 5.1 is a generalization of [St, (1)]. In proving [St, (1)], Stembridge uses the simple fact that if S is a tableau and $\xi = (\xi_1, \ldots, \xi_d) \in \mathbb{N}^d$, then $x^\xi x^S = x^{\xi+\omega(S)}$. The generalization of this fact which we will need in order to prove Lemma 5.1 is the following lemma. Define $(x\,|\,y)^\xi = (x_1\,|\,y)^{\xi_1} \cdots (x_d\,|\,y)^{\xi_d}$.

Lemma 7.1. *Let $R \in \mathcal{R}_{sub}(\mu)$ and let $\xi \in \mathbb{N}^d$. Then*

$$(x \mid y)^\xi (x \mid y)^R = \sum_{\substack{B \in \mathcal{B}_{sub}(\mu) \\ \widetilde{B} = R}} c_{\xi+1,B} \cdot (x \mid y)^{\xi+\omega(B^u)}.$$

In fact, we only need this lemma for $R \in \mathcal{R}(\mu)$. We prove this result more generally for $R \in \mathcal{R}_{sub}(\mu)$ only to allow for induction on the number of entries of R (and thus allow for the possibility that some boxes of R are empty). We remark that $\mathcal{R}_{\mathrm{sub}}(\mu)$ and $\mathcal{B}_{\mathrm{sub}}(\mu)$ were introduced in this paper solely to allow for induction in this proof.

Proof. By induction on the number of entries in R. Let a be an entry of R with value k, such that R has no entry of value k in any column to the left of a. Let α be the box containing a. Let $R' = R \setminus a$ be the the reverse subtableau which results from removing a from R.

If $B \in \mathcal{B}_{sub}(\mu)$ is such that $\widetilde{B} = R$, then the entry of B in box α, which we denote by B_α, must either be k or $\overline{k}$. Let B' denote $B \setminus B_\alpha$. The following three sets are in bijection with one another:

$$\begin{aligned}\{B \in \mathcal{B}_{sub}(\mu) \mid \widetilde{B} = R, B_\alpha = k\} &\longleftrightarrow \{B \in \mathcal{B}_{sub}(\mu) \mid \widetilde{B} = R, B_\alpha = \overline{k}\} \\ &\longleftrightarrow \{D \in \mathcal{B}_{sub}(\mu) \mid \widetilde{D} = R'\}.\end{aligned}$$

The first bijection simply adds a bar to B_α, and the second bijection removes B_α from B, mapping B to B'. For brevity, we denote $e_{B,\xi}(B_\alpha)$ and $f_B(B_\alpha)$ by just $e(B_\alpha)$ and $f(B_\alpha)$ respectively for the remainder of this proof. If B_α is unbarred, then

$$c_{\xi+1,B} = c_{\xi+1,B'} \quad \text{and} \quad (x \mid y)^{\xi+\omega(B^u)} = (x \mid y)^{\xi+\omega((B')^u)}(x_d - y_{e(B_\alpha)+1}).$$

On the other hand, if B_α is barred, then

$$c_{\xi+1,B} = c_{\xi+1,B'}(y_{e(B_\alpha)+1} - y_{f(B_\alpha)}) \quad \text{and} \quad (x \mid y)^{\xi+\omega(B^u)} = (x \mid y)^{\xi+\omega((B')^u)}.$$

Thus,

$$\begin{aligned}
&\sum_{\substack{B\in\mathcal{B}_{sub}(\mu)\\ \widetilde{B}=R}} c_{\xi+1,B}(x \mid y)^{\xi+\omega(B^u)} \\
&\qquad = \sum_{\substack{B\in\mathcal{B}_{sub}(\mu)\\ \widetilde{B}=R\\ B_\alpha=k}} c_{\xi+1,B}(x \mid y)^{\xi+\omega(B^u)} + \sum_{\substack{B\in\mathcal{B}_{sub}(\mu)\\ \widetilde{B}=R\\ B_\alpha=\overline{k}}} c_{\xi+1,B}(x \mid y)^{\xi+\omega(B^u)} \\
&\qquad = \sum_{\substack{B\in\mathcal{B}_{sub}(\mu)\\ \widetilde{B}=R\\ B_\alpha=k}} c_{\xi+1,B'}(x \mid y)^{\xi+\omega((B')^u)}(x_{B_\alpha} - y_{e(B_\alpha)+1}) \\
&\qquad\qquad + \sum_{\substack{B\in\mathcal{B}_{sub}(\mu)\\ \widetilde{B}=R\\ B_\alpha=\overline{k}}} c_{\xi+1,B'}(y_{e(B_\alpha)+1} - y_{f(B_\alpha)})(x \mid y)^{\xi+\omega((B')^u)} \\
&\qquad = \sum_{\substack{B\in\mathcal{B}_{sub}(\mu)\\ \widetilde{B}=R}} \Big(c_{\xi+1,B'}(x \mid y)^{\xi+\omega((B')^u)}(x_{B_\alpha} - y_{e(B_\alpha)+1}) \\
&\qquad\qquad + \quad c_{\xi+1,B'}(y_{e(B_\alpha)+1} - y_{f(B_\alpha)})(x \mid y)^{\xi+\omega((B')^u)}\Big) \\
&\qquad = \sum_{\substack{B\in\mathcal{B}_{sub}(\mu)\\ \widetilde{B}=R}} c_{\xi+1,B'}(x \mid y)^{\xi+\omega((B')^u)}(x_{B_\alpha} - y_{f(B_\alpha)}) \\
&\qquad = \sum_{\substack{D\in\mathcal{B}_{sub}(\mu)\\ \widetilde{D}=R'}} \Big(c_{\xi+1,D}(x \mid y)^{\xi+\omega(D^u)}\Big)(x_{B_\alpha} - y_{f(B_\alpha)}) \\
&\qquad = (x \mid y)^{\xi}(x \mid y)^{R'}(x_{B_\alpha} - y_{f(B_\alpha)}) \\
&\qquad = (x \mid y)^{\xi}(x \mid y)^{R}.
\end{aligned}$$

□

Proof of Lemma 5.1.

$$\begin{aligned}
a_{\lambda+\rho}(x\,|\,y)s_{\mu}(x\,|\,y) &\overset{(a)}{=} \sum_{\sigma\in S_d}\sum_{R\in\mathcal{R}(\mu)} \operatorname{sgn}(\sigma)(x \mid y)^{\sigma(\lambda+\rho)}(x \mid y)^{R} \\
&\overset{(b)}{=} \sum_{\sigma\in S_d}\sum_{R\in\mathcal{R}(\mu)} \sum_{\substack{B\in\mathcal{B}(\mu)\\ \widetilde{B}=R}} c_{\sigma(\lambda+\rho+1),B}\operatorname{sgn}(\sigma)(x \mid y)^{\sigma(\lambda+\rho)+\omega(B^u)} \\
&\overset{(c)}{=} \sum_{\sigma\in S_d}\sum_{R\in\mathcal{R}(\mu)} \sum_{\substack{B\in\mathcal{B}(\mu)\\ \widetilde{B}=R}} \sum_{\substack{H\in\mathcal{H}(\mu)\\ \overline{H}=B}} d_{\sigma(\lambda+\rho+1),H}\operatorname{sgn}(\sigma)(x \mid y)^{\sigma(\lambda+\rho)+\omega(H^u)} \\
&= \sum_{\sigma\in S_d}\sum_{H\in\mathcal{H}(\mu)} d_{\sigma(\lambda+\rho+1),H}\operatorname{sgn}(\sigma)(x \mid y)^{\sigma(\lambda+\rho)+\omega(H^u)} \\
&\overset{(d)}{=} \sum_{\sigma\in S_d}\sum_{H\in\mathcal{H}(\mu)} d_{\sigma(\lambda+\rho+1),\sigma H}\operatorname{sgn}(\sigma)(x \mid y)^{\sigma(\lambda+\rho)+\omega((\sigma H)^u)} \\
&\overset{(e)}{=} \sum_{\sigma\in S_d}\sum_{H\in\mathcal{H}(\mu)} d_{\lambda+\rho+1,H}\operatorname{sgn}(\sigma)(x \mid y)^{\sigma(\lambda+\rho+\omega(H^u))} \\
&= \sum_{\sigma\in S_d}\sum_{B\in\mathcal{B}(\mu)} \sum_{\substack{H\in\mathcal{H}(\mu)\\ \overline{H}=B}} d_{\lambda+\rho+1,H}\operatorname{sgn}(\sigma)(x \mid y)^{\sigma(\lambda+\rho+\omega(H^u))} \\
&\overset{(c)}{=} \sum_{\sigma\in S_d}\sum_{B\in\mathcal{B}(\mu)} c_{\lambda+\rho+1,B}\operatorname{sgn}(\sigma)(x \mid y)^{\sigma(\lambda+\rho+\omega(B^u))} \\
&\overset{(a)}{=} \sum_{B\in\mathcal{B}(\mu)} c_{\lambda+\rho+1,B}\,a_{\lambda+\rho+\omega(B^u)}(x\,|\,y) \overset{(f)}{=} \sum_{B\in\mathcal{B}(\mu)} c_{\lambda,B}\,a_{\lambda+\rho+\omega(B^u)}(x\,|\,y).
\end{aligned}$$

Equality (a) follows from the definition of a_μ, noting that $\sigma(\lambda+\rho)+1=\sigma(\lambda+\rho+1)$; (b) follows from Lemma 7.1, setting $S=R$ and $\xi=\sigma(\lambda+\rho)$; (c) from (8), with $\xi=\sigma(\lambda+\rho)$; (e) from (10); and (f) from (7). For (d), we use the fact that for a fixed σ and arbitrary decomposition $\sigma=\sigma_{i_1}\cdots\sigma_{i_t}$, since each s_{i_j} is an involution on $\mathcal{H}(\mu)$, as H runs over all elements of $\mathcal{H}(\mu)$, so does σH. $\square$

8. PROOF OF LEMMA 5.2

By (9), Lemma 5.2 is equivalent to the following lemma, whose statement and proof generalize arguments in [St]. For $H\in\mathcal{H}(\mu)$ and j a nonnegative integer, define $H_{<j}$ to be the sub-hatted tableau of H consisting of the portion of H lying in columns to the right of j, and $H^u_{<j}=(H_{<j})^u$ (and similarly for $H_{\le j}$, $H_{>j}$, etc.).

Lemma 8.1. *Let $\lambda\in\mathcal{P}_n$. Then*

$$\sum d_{\lambda+\rho+1,H}\,a_{\lambda+\rho+\omega(H^u)}(x\,|\,y)=0, \tag{11}$$

the sum being over all $H\in\mathcal{H}(\mu)$ for which $\lambda+\omega(H^u_{\le j})\notin\mathcal{P}_d$ for some j.

Proof. We call $H\in\mathcal{H}(\mu)$ for which $\lambda+\omega(H^u_{\le j})\notin\mathcal{P}_d$ for some j a *Bad Guy*. Let H be a Bad Guy, and let j be minimal such that $\lambda+\omega(H^u_{\le j})\notin\mathcal{P}_d$. Having selected j, let i be minimal such that $(\lambda+\omega(H^u_{\le j}))_i<(\lambda+\omega(H^u_{\le j}))_{i+1}$. Since $(\lambda+\omega(H^u_{\le j-1}))_i\ge(\lambda+\omega(H^u_{\le j-1}))_{i+1}$ (by the minimality of j), we must have $(\lambda+\omega(H^u_{\le j-1}))_i=(\lambda+\omega(H^u_{\le j-1}))_{i+1}$, and column j of H must have an unhatted

$i+1$ but not an unhatted i. Thus

$$(\lambda+\rho+1+\omega(H^u_{\leq j}))_i = (\lambda+\rho+1+\omega(H^u_{\leq j}))_{i+1}. \tag{12}$$

Define H^* to be the reverse tableau of shape μ obtained from H by replacing $H_{>j}$ by $s_i(H_{>j})$. Notice first that H^* is still semistandard. Indeed, since σ_i applied to $H_{>j}$ can only change the values of its entries from i to $i+1$ and visa-versa, the only possible violation of semistandardness of H^* would occur under the following scenario: (a) H has an entry a of value i in column j (which has to be either an $\widehat{\imath}$ or $\hat{\imath}$), (b) H has an entry b of value i immediately to the left of a, and (c) s_i applied to $H_{>j}$ changes the value of b to $i+1$. However, this scenario is impossible. If (a) and (b) both hold, then since H is semistandard, the entry of H immediately below b must have value $i+1$ (we remark that the *reverse* shape of the tableau is critical here). Therefore the entry in box b is not a free entry of $H_{>j}$, so s_i does not change its value, i.e., (c) is violated. Notice second that H^* is still a Bad Guy, since $H^*_{\leq j} = H_{\leq j}$. Thus $H \mapsto H^*$ gives an involution on the set of Bad Guys of $H(\mu)$.

We define maps $b_l^* : H^l \to (H^*)^l$ and $b_r^* : H^r \to (H^*)^r$, as follows. If $a \in (H_{\leq j})^l$, then define $b_l^*(a) = a$. If $a \in (H_{>j})^l$, then during the construction of H^*, in the process of applying s_i to $H_{>j}$, a is mapped to $b_l(a) \in (H_{>j})^l$. This same element $b_l(a)$, regarded as an element of $(H^*)^l$, is denoted by $b_l^*(a)$. The map b_r^* is defined analogously.

We wish to show that for $a \in H^l$,

$$e_{\lambda+\rho+1,H}(a) = e_{\lambda+\rho+1,H^*}(b_l^*(a)), \tag{13}$$

and for $a \in H^r$,

$$f_H(a) = f_{H^*}(b_r^*(a)). \tag{14}$$

For $a \in (H_{\leq j})^l$ or $a \in (H_{\leq j})^r$, both (13) and (14) are obvious. The proof of (14) for $a \in (H_{>j})^r$ follows in much the same manner as the proof of Lemma 6.4(v).

It remains to prove (13) for $a \in (H_{>j})^l$. For such a, by Lemma 6.4(iii),

$$\sigma_i\omega({}_{j<}H^u_{<c(a)}) = \omega(s_i({}_{j<}H^u_{<c(b_l(a))})) = \omega({}_{j<}(H^*)^u_{<c(b_l^*(a))}), \tag{15}$$

where ${}_{l<}H_{<m} := H_{>l} \cap H_{<m}$. By (12),

$$\sigma_i(\lambda+\rho+1+\omega(H^u_{\leq j})) = \lambda+\rho+1+\omega(H^u_{\leq j}) = \lambda+\rho+1+\omega((H^*)^u_{\leq j}). \tag{16}$$

Thus

$$\begin{aligned}
e_{\lambda+\rho+1,H}(a) &= \Big(\lambda+\rho+1+\omega(H^u_{<c(a)}))\Big)_{|a|} \\
&= \Big(\sigma_i(\lambda+\rho+1+\omega(H^u_{<c(a)}))\Big)_{\sigma_i|a|} \\
&\overset{(a)}{=} \Big(\sigma_i(\lambda+\rho+1+\omega(H^u_{<c(a)}))\Big)_{|b_l^*(a)|} \\
&= \Big(\sigma_i(\lambda+\rho+1+\omega({}_{j<}H^u_{<c(a)})+\omega(H^u_{\leq j}))\Big)_{|b_l^*(a)|} \\
&= \Big(\sigma_i(\omega({}_{j<}H^u_{<c(a)}))+\sigma_i(\lambda+\rho+1+\omega(H^u_{\leq j}))\Big)_{|b_l^*(a)|} \\
&\overset{(b)}{=} \Big(\omega({}_{j<}(H^*)^u_{<c(b_l^*(a))})+\lambda+\rho+1+\omega((H^*)^u_{\leq j})\Big)_{|b_l^*(a)|}
\end{aligned}$$

$$= \Big(\lambda+\rho+1+\omega((H^*)^u_{<c(b_l^*(a))})\Big)_{|b_l^*(a)|}$$
$$= e_{\lambda+\rho+1,H^*}(b_l^*(a)).$$

Equality (a) follows from Lemma 6.4(i); (b) follows from (15) and (16). This completes the proofs of (13) and (14).

Now (13) and (14) imply

$$\begin{aligned} d_{\lambda+\rho+1,H} &= \prod_{a\in H^l} y_{e_{\lambda+\rho+1,H}(a)} \prod_{a\in H^r}(-y_{f_H(a)}) \\ &= \prod_{a\in H^l} y_{e_{\lambda+\rho+1,H^*}(b_l^*(a))} \prod_{a\in H^r}(-y_{f_{H^*}(b_l^*(a))}) \\ &= \prod_{a\in (H^*)^l} y_{e_{\lambda+\rho+1,H^*}(a)} \prod_{a\in (H^*)^r}(-y_{f_{H^*}(a)}) = d_{\lambda+\rho+1,H^*}. \end{aligned} \tag{17}$$

By $\sigma_i\omega(H^u_{>j}) = \omega((H^*)^u_{>j})$ and (16), $\sigma_i(\lambda+\rho+\omega(H^u)) = (\lambda+\rho+\omega((H^*)^u))$; thus

$$a_{\lambda+\rho+\omega(H^u)}(x\,|\,y) = -a_{\lambda+\rho+\omega((H^*)^u)}(x\,|\,y). \tag{18}$$

By (17) and (18), the contributions to (11) of two Bad Guys paired under the involution $H \mapsto H^*$ are negatives, and thus cancel. If a Bad Guy is paired with itself under $H \mapsto H^*$, then (18) implies that its contribution to (11) is 0. $\square$

9. Bijection with Knutson-Tao Puzzles

In this section we give a weight-preserving bijection between positive equivariant Littlewood-Richardson skew tableaux and Knutson-Tao puzzles; thus both combinatorial objects compute identical expressions for the structure constants $c^\nu_{\lambda,\mu}$ and $C^\nu_{\lambda,\mu}$, $\lambda,\mu,\nu\in\mathcal{P}_{d,n}$. We begin by reviewing the construction of Knutson-Tao puzzles.

9.1. **Puzzles.** A **puzzle piece** is one of the eight figures shown in Figure 3, each of whose edges has length 1 unit. Each puzzle piece is either an equilateral triangles or a rhombus, together with a fixed orientation, and a labelling of each edge with either a 1 or 0. The rightmost puzzle piece in Figure 3 is called an **equivariant puzzle piece**; we color it cyan (or light gray on a black and white printer).



Figure 3. The eight puzzle pieces

A **puzzle** P is a partitioning of an equilateral triangle of side length n into puzzle pieces (see Figure 4). Implicit in this definition is that if two puzzle pieces of P share an edge, then both puzzle pieces must have the same label on that edge. The large equilateral triangle forming the boundary of P is called simply the **boundary** of P and denoted by ∂P. The northeast, northwest, and south sides of the boundary are denoted by ∂P_{NE}, ∂P_{NW}, and ∂P_{S} respectively. One forms three n-digit binary words by reading the labels along the three sides of ∂P: the labels of ∂P_{NE} are read from top to bottom, the labels of ∂P_{NW} from bottom to top, and the labels of ∂P_{S} from left to right. To these three binary words we associate three partitions of $\mathcal{P}_{d,n}$ under the map $w \mapsto (\eta_1, \ldots, \eta_d) \in \mathcal{P}_{d,n}$, where η_j is the number of zeros of w which lie to the right of the j-th one of w from the left (for example, $0110001010 \mapsto (5,5,2,1) \in \mathcal{P}_{4,10}$). Denote by $\mathcal{LP}^{\nu\,+}_{\lambda,\mu}$ the set of all puzzles P for which these three partitions are λ, μ, and ν, in that order.



FIGURE 4. A puzzle P, with $n = 9$, $d = 3$. The n-digit binary words of the NE, NW, and S sides of the boundary are 001001100, 001010010, and 101000100 respectively. Thus $P \in \mathcal{LP}^{\nu\,+}_{\lambda,\mu}$, where $\lambda = (4,2,2)$, $\mu = (4,3,1)$, and $\nu = (6,5,2)$.

For any equivariant puzzle piece of a puzzle P, draw two lines from the center of the puzzle piece to ∂P_{S}: one line L_1 parallel to ∂P_{NW} and the other L_2 parallel to ∂P_{NE} (see Figure 5). The line segment ∂P_{S} consists of n edges of puzzle pieces, which we number $1, 2, \ldots, n$ from right to left. The lines L_1 and L_2 cross ∂P_{S} in the center of two edges e and f respectively, where $e > f$. The **factorial weight of the puzzle piece** is $y_e - y_f$, and the **equivariant weight of the puzzle piece** is $Y_{n+1-f} - Y_{n+1-e}$. Let c_P denote the product of the factorial weights of all the equivariant puzzle pieces of P and C_P the product of the equivariant weights of all the equivariant puzzle pieces of P. For example, in Figure 4, $c_P = (y_8 - y_3)(y_3 - y_2)(y_3 - y_1)$ and $C_P = (Y_7 - Y_2)(Y_8 - Y_7)(Y_9 - Y_7)$.

Proposition 9.2. *There is a weight preserving bijection* $\Phi : \mathcal{LP}^{\nu\,+}_{\lambda,\mu} \to \mathcal{LR}^{\nu\,+}_{\lambda,\mu}$. *By weight-preserving, we mean that for* $P \in \mathcal{LP}^{\nu\,+}_{\lambda,\mu}$, c_P *and* $c_{\Phi(P)}$ *are equal, and moreover are identical expressions; and similarly for* C_P *and* $C_{\Phi(P)}$.

Proof. The bijection Φ, illustrated in Figure 6, generalizes Tao's 'proof without words' of the bijection between puzzles and tableaux in the nonequivariant setting



FIGURE 5. The equivariant puzzle piece has factorial weight $y_6 - y_3$ and equivariant weight $Y_5 - Y_2$.

[V, Figure 11]. The large triangle in the center of Figure 6 represents a generic puzzle P. The equivariant Littlewood-Richardson skew tableau $\Phi(P)$ is the skew barred tableau formed by placing the Young tableau on the top-right of Figure 6 above and to the right of the reverse barred tableau on the top left of the figure.

We list three properties of any $L \in \mathcal{LR}^{\nu\,+}_{\lambda,\mu}$:

(a) $L|_\mu$ is column strict;
(b) the unbarred column word of L is Yamanouchi; and
(c) $c_L > 0$.

Let $P \in \mathcal{LP}^{\nu\,+}_{\lambda,\mu}$. For $i \in \{1, \ldots, d\}$ (where $d = 4$ in Figure 6), there is a path P_i consisting of 1-triangles and rhombi which begins on ∂P_{NE}, moves only west or southwest, and ends on ∂P_{S} (see Figure 7). Each path P_i has **segments** $P_{i,j}$ consisting of the consecutive rhombi lying to the right of an upward pointing 1-triangle and to the left of either a downward pointing 1 triangle or ∂P_{NE}. We list three properties of P:

(a)' for $i = 2, \ldots, d$ and all j, the distance from the leftmost edge of $P_{i,j}$ to ∂P_{NE} is greater than or equal to the distance from the leftmost edge of $P_{i-1,j}$ to to ∂P_{NE};
(b)' the interiors of the P_i do not touch; and
(c)' the interiors of all equivariant puzzle pieces lie above ∂P_{S}.

Given any $P \in \mathcal{LP}^{\nu\,+}_{\lambda,\mu}$, Figure 6 shows how to construct a skew barred tableau $\Phi(P)$. Properties (a)', (b)', and (c)' of P imply properties (a), (b), and (c) of $\Phi(P)$ respectively. Conversely, given any $L \in \mathcal{LR}^{\nu\,+}_{\lambda,\mu}$, Figure 6 shows how to construct a puzzle $\Phi^{-1}(L)$. Properties (a), (b), and (c) of L ensure that the puzzle $\Phi^{-1}(L)$ can be constructed, and imply that it satisfies (a)', (b)', and (c)' respectively. Uniqueness is clear.

To each equivariant puzzle piece of P there corresponds a barred entry of $\Phi(P)$, and they both determine the same factor $y_i - y_j$ of c_P and $c_{\Phi(P)}$ respectively. Therefore Φ is weight preserving. □

Using Corollary 2.8 and Proposition 9.2, we obtain a new proof of the following theorem, which is due to Knutson and Tao [KT].

Theorem 9.3 (Knutson-Tao). $c^\nu_{\lambda,\mu} = \sum_{P \in \mathcal{LP}^{\nu\,+}_{\lambda,\mu}} c_P$ *and* $C^\nu_{\lambda,\mu} = \sum_{P \in \mathcal{LP}^{\nu\,+}_{\lambda,\mu}} C_P$.



Figure 6. A generic puzzle P (center), and its associated positive equivariant Littlewood-Richardson skew tableau $\Phi(P)$ (top-right, top-left). In P, black represents regions of 1 triangles, green (dark gray) represents regions of 0 triangles, white represents regions of non-equivariant rhombi, and cyan (light gray) represents regions of equivariant rhombi.



FIGURE 7. The paths P_i, $i = 1, \ldots, 4$, of the puzzle P of Figure 6. The segments $P_{i,j}$ of each path are shaded. The segments may contain two types of puzzle pieces: equivariant puzzle pieces and rhombi with horizontal 0-edges.



Figure 8. The puzzle P of Figure 4 is redrawn using the coloring scheme described in Figure 6. All edge labels other than those on the boundary of the puzzle are suppressed. $\Phi(P)$ is also shown.



Figure 9. All elements of $\mathcal{LP}^{\nu\,+}_{\lambda,\mu}$ and $\mathcal{LR}^{\nu\,+}_{\lambda,\mu}$, for $d = 2$, $n = 4$, $\lambda = (1, 1)$, $\mu = (2, 1)$, and $\nu = (2, 1)$. We have $c^{\mu}_{\lambda,\mu} = (y_4 - y_3)(y_2 - y_1) + (y_3 - y_1)(y_2 - y_1)$ and $C^{\mu}_{\lambda,\mu} = (Y_2 - Y_1)(Y_4 - Y_3) + (Y_4 - Y_2)(Y_4 - Y_3)$.

9.4. **Trapezoid Puzzles.** We next extend Φ to a bijection onto $\mathcal{LR}^{\nu}_{\lambda,\mu}$. To do so, we increase the size of the domain of Φ by defining generalizations of puzzles, which we call *trapezoid puzzles*. The extention of Φ, which we also denote by Φ, allows us to view nonnegativity from the point of view of trapezoid puzzles rather than equivariant Littlewood-Richardson skew tableaux.

Consider the isosceles trapezoid T formed by placing an equilateral triangle of side length n on top of a rhombus of side length n (see Figure 10). The boundary of T, denoted by of ∂T, is divided into 5 parts: northeast, northwest, east, west, and south (denoted by ∂T_{NE}, ∂T_{NW}, ∂T_{E}, ∂T_{W}, and ∂T_{S}). The northeast and northwest boundaries of T are the northeast and northwest boundaries of the equilateral triangle, and the east, west, and south boundaries of T are the east, west, and south boundaries of the rhombus. A **trapezoid puzzle** is a partitioning of T into puzzle pieces in such a way that all labels of ∂T_{E} and ∂T_{W} are 0's. Denote by $\mathcal{LP}^{\nu}_{\lambda,\mu}$ the set of all trapezoid puzzles whose n-digit binary words read from ∂T_{NE}, ∂T_{NW}, and ∂T_{S} correspond to partitions λ, μ, and ν respectively.

Let D denote the line segment forming the south border of the triangle (and the north border of the rhombus). For any equivariant puzzle piece of a trapezoid puzzle P, draw two lines from the center of the puzzle piece to D: one line L_1 parallel to ∂T_{NW} and the other L_2 parallel to ∂T_{NE}. The lines L_1 and L_2 cross D at $e - .5$ and $f - .5$ units from its right endpoint, respectively (e, f are both integers). If the equivariant puzzle piece lies above D, then $e > f$; if it lies below D, then $e < f$; if it is bisected by D, then $e = f$. The **factorial weight of the puzzle piece** is $y_e - y_f$, and the **equivariant weight of the puzzle piece** is $Y_{n+1-f} - Y_{n+1-e}$. Let c_P denote the product of the factorial weights of all the equivariant puzzle pieces of P and C_P the product of the equivariant weights of all the equivariant puzzle pieces of P.

A puzzle can be viewed as a trapezoid puzzle all of whose 1-triangles lie above D. In this way $\mathcal{LP}^{\nu\,+}_{\lambda,\mu}$ may be viewed as a subset of $\mathcal{LP}^{\nu}_{\lambda,\mu}$, and the inclusion is weight-preserving. A diagram very similar to Figure 6, but for trapezoid puzzles instead of puzzles, can be used to prove

Proposition 9.5. *The bijection* $\Phi : \mathcal{LP}^{\nu\,+}_{\lambda,\mu} \to \mathcal{LR}^{\nu\,+}_{\lambda,\mu}$ *of Proposition 9.2 extends to a weight preserving bijection* $\Phi : \mathcal{LP}^{\nu}_{\lambda,\mu} \to \mathcal{LR}^{\nu}_{\lambda,\mu}$.

One makes the following two observations: if $P \in \mathcal{LP}^{\nu}_{\lambda,\mu} \setminus \mathcal{LP}^{\nu\,+}_{\lambda,\mu}$, i.e., if P has a 1-triangle lying below D, then (i) at least one equivariant puzzle piece must be bisected by D, and (ii) stronger, at least one equivariant puzzle piece corresponding to a bottom row element of $\Phi(P)$ must be bisected by D. Although we have not included a diagram such as Figure 6 for trapezoid puzzles, these two statements can nevertheless be seen in Figure 6 itself, if one imagines placing the required side length n rhombus underneath the figure, and 'stretching' the 'paths', expanding without breaking the 'loops', so that the 1-triangles are pushed into the rhombus below. Statement (i) implies that $c_P = C_P = 0$. Combined with Proposition 9.5, it gives a simpler proof of Proposition 2.7. Statement (ii) proves that Lemma 4.2(iii) implies Lemma 4.2(i). That Lemma 4.2(i) implies Lemma 4.2(ii) can also be seen easily by considering trapezoid puzzles (or puzzles).

By Theorem 9.3 and Proposition 9.5, we have

Corollary 9.6. $c^{\nu}_{\lambda,\mu} = \displaystyle\sum_{P \in \mathcal{LP}^{\nu}_{\lambda,\mu}} c_P$ *and* $C^{\nu}_{\lambda,\mu} = \displaystyle\sum_{P \in \mathcal{LP}^{\nu}_{\lambda,\mu}} C_P$.


1
1 1 $\overline{1}$ $\overline{2}$ 2
1 $\overline{2}$ 2 $\overline{3}$ 3 $\overline{3}$ $\overline{3}$ $\overline{3}$
1 1 1 1 1
2 2
3
1 1 1 1 1 1 1 1 1
2 2 2 2
3 3
0 1 0 0 0 0 0 1 0 0 1 0 0

FIGURE 10. A trapezoid puzzle P (center), for $d = 3$, $n = 13$, $\lambda = (5, 2, 1)$, $\mu = (8, 5, 1)$, $\nu = (9, 4, 2)$; and the corresponding equivariant Littlewood-Richardson skew tableau $\Phi(P)$ (top-right, top-left). The line D separating the triangle from the rhombus is darkened. The fact that 1-triangles lie below D implies that $c_P = C_P = 0$. Indeed, $c_P = (y_9 - y_4)(y_5 - y_2)(y_2 - y_1)(y_2 - y_2)(y_2 - y_3)(y_3 - y_5)(y_6 - y_8) = 0$, and $C_P = (Y_{10} - Y_5)(Y_{12} - Y_9)(Y_{13} - Y_{12})(Y_{12} - Y_{12})(Y_{11} - Y_{12})(Y_9 - Y_{11})(Y_6 - Y_8) = 0$.

REFERENCES

[Bi] S. C. Billey, *Kostant polynomials and the cohomology ring for G/B*, Duke Math. J. **96** (1999), no. 1, 205–224.

[BK] E. A. Bender and D. E. Knuth, *Enumeration of plane partitions*, J. Combinatorial Theory Ser. A **13** (1972), 40–54.

[BL] L. C. Biedenharn and J. D. Louck, *A new class of symmetric polynomials defined in terms of tableaux*, Adv. in Appl. Math. **10** (1989), no. 4, 396–438.

[CL] W. Y. C. Chen and J. D. Louck, *The factorial Schur function*, J. Math. Phys. **34** (1993), no. 9, 4144–4160.

[F1] W. Fulton, *Young tableaux*, London Mathematical Society Student Texts, vol. 35, Cambridge University Press, Cambridge, 1997.

[F2] W. Fulton, *Equivariant cohomology in algebraic geometry*, Eilenberg lectures, Columbia University, 2007. Available at `http://www.math.lsa.umich.edu/dandersn/eilenberg`.

[GG] I. Goulden and C. Greene, *A new tableau representation for supersymmetric Schur functions*, J. Algebra **170** (1994), no. 2, 687–703.

[Gr] W. Graham, *Positivity in equivariant Schubert calculus*, Duke Math. J. **109** (2001), no. 3, 599–614.

[IN] T. Ikeda and H. Naruse, *Excited Young diagrams and equivariant Schubert calculus*, arXiv:math.AG/0703637.

[Kr] V. Kreiman, *Schubert classes in the equivariant K-theory and equivariant cohomology of the Grassmannian*, arXiv:math.AG/0512204.

[KT] A. Knutson and T. Tao, *Puzzles and (equivariant) cohomology of Grassmannians*, Duke Math. J. **119** (2003), no. 2, 221–260.

[KTW] A. Knutson, T. Tao, and C. Woodward, *The honeycomb model of* $\mathrm{GL}_n(\mathbb{C})$ *tensor products. II. Puzzles determine facets of the Littlewood-Richardson cone*, J. Amer. Math. Soc. **17** (2004), no. 1, 19–48 (electronic).

[La] A. Lascoux, *Puissances extérieures, déterminants et cycles de Schubert*, Bull. Soc. Math. France **102** (1974), 161–179.

[LS1] A. Lascoux and M.-P. Schützenberger, *Polynômes de Schubert*, C. R. Acad. Sci. Paris Sér. I Math. **294** (1982), no. 13, 447–450.

[LS2] ———, *Structure de Hopf de l'anneau de cohomologie et de l'anneau de Grothendieck d'une variété de drapeaux*, C. R. Acad. Sci. Paris Sér. I Math. **295** (1982), no. 11, 629–633.

[LR] D. E. Littlewood and A. R. Richardson, *Group characters and algebra*, Philos. Trans. Roy. Soc. London Ser. A **233** (1934).

[Ma1] I. G. Macdonald, *Schur functions: theme and variations*, Séminaire Lotharingien de Combinatoire (Saint-Nabor, 1992), Publ. Inst. Rech. Math. Av., vol. 498, Univ. Louis Pasteur, Strasbourg, 1992, pp. 5–39.

[Ma2] ———, *Symmetric functions and Hall polynomials*, second ed., Oxford Mathematical Monographs, The Clarendon Press Oxford University Press, New York, 1995, With contributions by A. Zelevinsky, Oxford Science Publications.

[Mc] P. J. McNamara, *Factorial Grothendieck polynomials*, Electron. J. Combin. **13** (2006), no. 1, Research Paper 71, 40 pp. (electronic).

[Mi] L. Mihalcea, *Giambelli formulae for the equivariant quantum cohomology of the Grassmannian*, arXiv:math.CO/0506335.

[Mo1] A. I. Molev, *Littlewood-Richardson polynomials*, arXiv:0704.0065.

[Mo2] A. I. Molev, *Factorial supersymmetric Schur functions and super Capelli identities*, Kirillov's seminar on representation theory, Amer. Math. Soc. Transl. Ser. 2, vol. 181, Amer. Math. Soc., Providence, RI, 1998, pp. 109–137.

[MS] A. I. Molev and B. E. Sagan, *A Littlewood-Richardson rule for factorial Schur functions*, Trans. Amer. Math. Soc. **351** (1999), no. 11, 4429–4443.

[R] S. Robinson, *A Pieri-type formula for* $H^*_T(\mathrm{SL}_n(\mathbb{C})/B)$, J. Algebra **249** (2002), no. 1, 38–58.

[Sa] B. E. Sagan, *The symmetric group*, second ed., Graduate Texts in Mathematics, vol. 203, Springer-Verlag, New York, 2001, Representations, combinatorial algorithms, and symmetric functions.

[St] J. R. Stembridge, *A concise proof of the Littlewood-Richardson rule*, Electron. J. Combin. **9** (2002), no. 1, Note 5, 4 pp. (electronic).

[V] R. Vakil, *A geometric Littlewood-Richardson rule*, Ann. of Math. (2) **164** (2006), no. 2, 371–421, Appendix A written with A. Knutson.

[Z] A. V. Zelevinsky, *A generalization of the Littlewood-Richardson rule and the Robinson-Schensted-Knuth correspondence*, J. Algebra **69** (1981), no. 1, 82–94.

Department of Mathematics, University of Georgia, Athens, GA 30602 USA
E-mail address: `vkreiman@math.uga.edu`